

DRSABCD during COVID-19



D

DANGER

- Check scene for any dangers
- Don PPE, following the correct process

**DO NOT
ATTEND PATIENT
WITHOUT PPE**

R

RESPONSE

Is the patient conscious?

- Assess response from **1.5m away**
- If no response, approach and check response to touch and talk

S

SEND FOR HELP

- Call 000/SurfCom for assistance
- Inform service you are wearing PPE

A

AIRWAY

- Assess airway in the position the patient is found
- If required, roll onto side and allow fluids to drain without assistance

**DO NOT USE
SUCTION**

**DO NOT USE
OP AIRWAYS**

B

BREATHING

- Apply oxygen via a therapy mask at 8LPM. Do not use more than 8LPM

DO NOT USE BVM

**DO NOT GIVE
RESCUE BREATHS**

C

COMPRESSIONS

- Start compressions at a rate of 100-120 per minute
- Compress at least 1/3 of chest

D

DEFIBRILLATION

Early defibrillation saves lives

- Attach AED and follow prompts

SAFE

SAFELY REMOVE PPE

The risk of contamination is highest during the removal of PPE

- Carefully remove PPE following the correct order and process
- Have a buddy watching you and ensure you do not touch your face

**DO NOT COMPLETE
REPORTS WHILE
WEARING PPE**

STAY COVID SAFE

For more information visit sls.com.au/covid-19/